



SULFONYL FLUORIDES (-SO₂F): MORE OPTIONS FOR DRUG DESIGN


Sulfonyl chlorides (-SO₂Cl) are widely used in medicinal chemistry and agrochemistry as precursors to pharmacologically important sulfonamides. Many sulfonyl chlorides with heteroaromatic substituents, however, are unstable due to SO₂ extrusion. More stable sulfonyl fluorides (-SO₂F) in many cases are the only option to synthesize the desired sulfonamides. They are less reactive, so that they might even have a free aliphatic amino groups in their structure. Besides unique monofunctional sulfonyl fluorides, *Enamine* offers a wide array of scaffolds and linker compounds.



Properties of sulfonyl fluorides




-SO₂F probes in chemical biology



The SO₂F group covalently binds to the residues of serine, threonine, tyrosine, lysine, cysteine, and histidine in proteins. Sulfonyl fluorides are widely used as chemical probes and covalent protein inhibitors.

Our offer: >200 Sulfonyl fluorides (-SO₂F) in gram amounts in stock
Custom synthesis of further analogues and compound libraries



References

¹ A. Narayanan *et al. Chem. Sci.* **2015**, 2650.
² J. Dong *et al. Angew. Chem. Int. Ed.* **2014**, 9430.

³ A. Garcia-Rubia *et al. Angew. Chem. Int. Ed.* **2011**, 10927.
⁴ S. W. Wright *et al. J. Org. Chem.* **2006**, 1080.

⁵ S. Caddick *et al. Org. Lett.* **2002**, 2549.



Search & Buy on-line at EnamineStore.com
Look for more at Chem-Space.com

BB@enamine.net, www.enamine.net